

Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at http://about.jstor.org/participate-jstor/individuals/early-journal-content.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

1065 June 28, 1912

MORTALITY—FOREIGN AND INSULAR—COUNTRIES AND CITIES (Untabulated).

Austria-Hungary—Brünn.—Month of April, 1912. Population, 108,944. Total number of deaths from all causes 310, including diphtheria 3, measles 14, scarlet fever 2, tuberculosis 69.

Brazil—Ceara.—Month of April, 1912. Population, 60,000. Total number of deaths from all causes 195, including tuberculosis 20, typhoid fever 5. There were 2 deaths from yellow fever.

British East Africa—Mombasa.—Month of April, 1912. Population, 26,000. Total number of deaths from all causes 45, including smallpox 1, tuberculosis 2, typhoid fever 1. Cases reported: Typhoid fever 1.

Canada—Hamilton.—Month of May, 1912. Population, 80,000. Total number of deaths from all causes 121, including diphtheria 1, measles 5, scarlet fever 1, tuberculosis 7. Cases reported: Diphtheria 13, measles 466, scarlet fever 20, tuberculosis 3.

Great Britain.—Week ended May 25, 1912.

England and Wales.—The deaths registered in 95 great towns correspond to an annual rate of 13 per 1,000 of the population, which is estimated at 17,639,816.

Ireland.—The deaths registered in 22 principal town districts correspond to an annual rate of 16.9 per 1,000 of the population, which is estimated at 1,157,014. The lowest rate was recorded at Clonmel, viz, 00.0, and the highest at Newry, viz, 26.2 per 1,000.

Scotland.—The deaths registered in 18 principal towns correspond to an annual rate of 15.4 per 1,000 of the population, which was estimated at 2,182,400. The lowest rate was recorded at Clydebank, viz, 6.5, and the highest at Ayr, viz, 21.8 per 1,000. The total number of deaths from all causes was 644, including diphtheria 6, measles 29, scarlet fever 3, typhoid fever 2.

Guam.—Three weeks ended April 19, 1912. Population, 9,000. Total number of deaths from all causes 16, including tuberculosis 4. There were 69 cases of hookworm disease.

Hawaii—*Hilo*.—Four weeks ended May 25, 1912. Population, 3,500; total number of deaths from all causes 4, including diphtheria 2, smallpox 1, tuberculosis 2. Cases reported: Diphtheria 3, tuberculosis 3.

ISLE OF PINES.—Nucva Gerona.—Twenty days ended June 10, 1912. Population, 892. Total number of deaths from all causes 1, from tuberculosis.

New Zealand.—Month of March, 1912.

Auckland.—Population, 104,728. Total number of deaths 64, including tuberculosis 1, typhoid fever 1.

June 28, 1912 1066

Christchurch.—Population, 82,004. Total number of deaths 46. Dunedin.—Population, 65,690. Total number of deaths 42, including tuberculosis 1.

Wellington.—Population, 71,427. Total number of deaths 48.

South Africa—Bloemfontein.—Month of April, 1912. Population, 22,000. Total number of deaths from all causes 42, including tuberculosis 2, typhoid fever 2. Cases reported: Scarlet fever 5, tuberculosis 3, typhoid fever 7. Remarks: Of the above, 28 deaths from all causes and 2 from tuberculosis; 1 case of typhoid fever and three cases of tuberculosis were reported from the colored population only.

Spain—Malaga.—Month of March, 1912. Population, 135,000. Total number of deaths from all causes 346, including diphtheria 2, measles 8, smallpox 5, tuberculosis 34, typhoid fever 1.

URUGUAY—Montevideo.—Month of March, 1912. Population, 343,-849. Total number of deaths from all causes 489, including diphtheria 1, tuberculosis 73, typhoid fever 12. Cases reported: Diphtheria 11, measles 7, scarlet fever 9, smallpox 1, tuberculosis 115, typhoid fever 60.

By authority of the Secretary of the Treasury:

Rupert Blue, Surgeon General, United States Public Health and Marine-Hospital Service.

ADDITIONAL COPIES of this publication may be procured from the SUPERINTEND-ENT OF DOCUMENTS, Government Printing Office, Washington, D. C., at 5 cents per copy.

Subscription price, per volume - - - \$2

